

KUNS-1560
hep-lat/9902022

Low energy effective action of domain-wall fermion and the Ginsparg-Wilson relation

Yoshio Kikukawa* and Tatsuya Noguchi†
Department of Physics, Kyoto University
Kyoto 606-8502, Japan

February, 1999

Abstract

We derive the effective action of the light fermion field of the domain-wall fermion, which is referred as $q(x)$ by Furman and Shamir. The inverse of the effective Dirac operator turns out to be identical to the inverse of the truncated overlap Dirac operator, except a local contact term which would give the chiral symmetry breaking in the Ginsparg-Wilson relation. This result allows us to relate the light fermion field and the fermion field described by the truncated overlap Dirac operator and to understand the chiral property of the light fermion through the exact chiral symmetry based on the Ginsparg-Wilson relation.

*e-mail address: kikukawa@gauge.scphys.kyoto-u.ac.jp

†e-mail address: noguchi@gauge.scphys.kyoto-u.ac.jp

1 Introduction

Recently, our understanding of chiral symmetry on the lattice has substantially improved. Lattice Dirac operators have been obtained [1, 2], which are gauge covariant, define local actions [3] and satisfy the Ginsparg-Wilson relation [4]. The Ginsparg-Wilson relation

$$\gamma_5 D + D \gamma_5 = a D R \gamma_5 D \quad (1.1)$$

implies the exact chiral symmetry of the action under the transformation [5]

$$\delta\psi(x) = \gamma_5 (1 - a R D) \psi(x), \quad \delta\bar{\psi}(x) = \bar{\psi}(x) \gamma_5. \quad (1.2)$$

The explicit gauge covariant solution of the Ginsparg-Wilson relation has been derived by Neuberger [1] from the overlap formulation of chiral determinant [6]. It is defined through the hermitian Wilson-Dirac operator H with a negative mass in a certain range ¹,

$$D = \frac{1}{2a} \left(1 + \gamma_5 \frac{H}{\sqrt{H^2}} \right). \quad (1.3)$$

The locality properties of the Dirac operator has been examined by Hernández, Jansen and Lüscher [3]. The issue of the practical implementation of the Dirac operator has been studied by Neuberger [7], Edwards, Heller and Narayanan [8], Chiu [9] and A. Borici [10].

The domain-wall fermion [11] is the basis of the overlap Dirac operator. In a simplified formulation [12, 13, 14], the domain-wall fermion consists of N -flavor Wilson fermions with a certain flavor-mixing mass matrix. Due to its structure of the chiral hopping and the boundary condition in the flavor space, a single light Dirac fermion can emerge in the spectrum. This light fermion can be probed most suitably by the field variables at the boundary in the flavor space, which are referred as $q(x)$ and $\bar{q}(x)$ by Furman and Shamir. It has been argued that the chiral symmetry of the light fermion is preserved up to corrections suppressed exponentially in the number of flavors [13]. This fact has been observed numerically [15, 16] and have been found useful for the numerical simulation of lattice QCD [17, 18, 19]. The perturbative studies are found in [20, 21]. The authors refer the reader to [22] for recent review. In this context, the chiral transformation of this light fermion is defined as

$$\delta q(x) = \gamma_5 q(x), \quad \delta \bar{q}(x) = \bar{q}(x) \gamma_5. \quad (1.4)$$

¹We adopt the definition of the overlap Dirac operator so that the normalization of the factor one half is included. This leads to the Ginsparg-Wilson relation with $R = 2$.

The goal of this paper is to understand the chiral property of the light fermion of the domain-wall fermion from the point of view of the exact chiral symmetry based on the Ginsparg-Wilson relation. Several authors have discussed the direct relation between the domain-wall fermion and the Dirac fermion which is described by the overlap Dirac operator. Vranas has shown that in order for the subtraction of the massive modes of the domain-wall fermion, it is suitable to introduce the N -flavor Wilson-Dirac boson with the flavor-mixing mass matrix which is anti-periodic in the flavor space [23]. Neuberger has shown in [14] through the explicit calculation of the partition function how the overlap Dirac operator emerges in the limit of infinite number of flavors: the subtracted partition function at a finite flavor N can be written as a single determinant of the truncated overlap Dirac operator

$$D_N = \frac{1}{2a} \left(1 + \gamma_5 \tanh \frac{N}{2} a_5 \tilde{H} \right). \quad (1.5)$$

Note that $\tilde{H}$ here is defined through the transfer matrix of the five-dimensional Wilson fermion with a negative mass:

$$\tilde{H} = -\frac{1}{a_5} \ln T. \quad (1.6)$$

In the limit of infinite flavors, this reduces to the overlap Dirac operator Eq. (1.3), in which the hermitian Wilson-Dirac operator H is replaced by $\tilde{H}$. Thus the domain-wall fermion with the subtraction of the Pauli-Villars field is equivalent to the Dirac fermion described by the truncated overlap Dirac operator.

The relation between the light fermion field $q(x)$ and $\bar{q}(x)$ of the domain-wall fermion and the fermion field $\psi(x)$ and $\bar{\psi}(x)$ described by the overlap Dirac operator has also been suggested by several authors. Neuberger pointed out in [14] the correspondence between the mass term for $q(x)$

$$m \bar{q}(x) q(x) \quad (1.7)$$

and the mass term for $\psi(x)$

$$m \bar{\psi}(x) \left(1 - \frac{a}{2} RD \right) \psi(x). \quad (1.8)$$

It has also been noticed by the authors of [24, 25] that the transformation property of the operator $\left(1 - \frac{a}{2} RD \right) \psi(x)$ under the chiral transformation Eq. (1.2) is same as $q(x)$:

$$\delta \left\{ \left(1 - \frac{a}{2} RD \right) \psi(x) \right\} = \gamma_5 \left\{ \left(1 - \frac{a}{2} RD \right) \psi(x) \right\}. \quad (1.9)$$

These facts suggest that there could be a correspondence as

$$q(x), \quad \bar{q}(x) \iff \left(1 - \frac{a}{2}RD\right) \psi(x), \quad \bar{\psi}(x) \quad (1.10)$$

at least in the limit of infinite flavors.²

In this paper, we will further examine the above correspondence between the light fermion field of the domain-wall fermion and the Dirac field described by the (truncated) overlap Dirac operator. For this purpose, we derive the low energy effective action of the light fermion field by integrating out $N - 1$ heavy flavors of the domain-wall fermion:

$$S_N^{\text{eff}} = a^4 \sum_x \bar{q}(x) D_N^{\text{eff}} q(x). \quad (1.11)$$

As easily understood, this can be achieved by calculating the propagator of $q(x)$ and $\bar{q}(x)$, because it should be given by the inverse of the effective Dirac operator of these fields.³

$$\langle q(x) \bar{q}(y) \rangle = \frac{1}{a^4} D_N^{\text{eff}^{-1}}(x, y). \quad (1.12)$$

It turns out that the propagator of the light fermion is closely related to the inverse of the truncated overlap Dirac operator as follows:

$$\frac{a}{a_5} D_N^{\text{eff}^{-1}} + a\delta(x, y) = D_N^{-1}. \quad (1.13)$$

Namely, the inverse of the effective Dirac operator gives the inverse of the truncated overlap Dirac operator up to a local contact term. This contact term just takes account of the chiral symmetry breaking in the Ginsparg-Wilson relation, which holds true for the overlap Dirac operator in the limit of the infinite flavors.

The above relation allows us to relate the field variables of the light fermion, $q(x)$ and $\bar{q}(x)$, with the field variables described by the truncated overlap Dirac operator, $\psi(x)$ and $\bar{\psi}(x)$. Then we can clarify the relation between the almost preserved chiral symmetry of the domain-wall fermion under the transformation Eq. (1.4) and the (would-be) exact chiral symmetry of the Dirac fermion described by the (truncated) overlap Dirac operator

²This correspondence could be different if we adopt a different transformation of the exact chiral symmetry of Lüscher from Eq. (1.2).

³ In this paper, the bra-ket symbol $\langle \cdots \rangle$ denotes the Wick contraction of the fermion fields in it by their propagators, not the fermionic vacuum expectation values which must includes the weight of the fermion action. The bra-ket symbol with the subscript c denotes the connected contraction.

under the transformation Eq. (1.2). It is also possible to relate the low energy observables of the domain-wall fermion which are written in terms of $q(x)$ and $\bar{q}(x)$ to those written in terms of $\psi(x)$ and $\bar{\psi}(x)$

$$\mathcal{O}_{\text{DW}}[q, \bar{q}] = \mathcal{O}_N[\psi, \bar{\psi}; D_N] \quad (1.14)$$

and to examine the chiral properties of the observables through the (would-be) exact chiral symmetry based on the Ginsparg-Wilson relation.

This article is organized as follows. In section 2, we will derive the effective action of the light fermion field of the domain-wall fermion by integrating out the heavy $N - 1$ flavors. The calculation of the effective action (the propagator of the light fermion) is a straightforward application of that given by Neuberger in [14]. In section 3, we will establish the relation between the light fermion field and the fermion field described by the truncated overlap Dirac operator. Then we discuss the chiral properties of the light fermion from the point of view of the exact chiral symmetry based on the Ginsparg-Wilson relation. In section 4, we reexamine the axial anomaly of the domain-wall fermion [26] in view of the axial anomaly associated with the exact chiral symmetry [27, 5, 28]. In section 5, the contribution of the Pauli-Villars field to the currents and the axial anomaly is examined.

2 Low energy effective action of the domain-wall fermion

We first review briefly the domain-wall fermion and its relation to the Dirac fermion described by the overlap Dirac operator. Then we evaluate the low energy effective action of the light fermion of the domain-wall fermion and discuss the result in relation to the Ginsparg-Wilson relation.

2.1 Light fermion of the domain-wall fermion

The domain-wall fermion, in its simplified formulation [12, 13, 14], consists of N -flavor Wilson fermions⁴

$$S_{\text{DW}} = \sum_{s,t=1}^N a^4 \sum_x \bar{\psi}_s(x) \left\{ \gamma_\mu \frac{1}{2} (\nabla_\mu + \nabla_\mu^*) \delta_{st} + P_L M_{st} + P_R M_{st}^\dagger \right\} \psi_t(x) \quad (2.1)$$

⁴In this paper, we assume that the number of flavor N is even.

with a certain flavor-mixing mass matrix: in the case with $N = 6$, it is given by

$$M_{st} = \frac{1}{a_5} \begin{pmatrix} B & -1 & 0 & 0 & 0 & 0 \\ 0 & B & -1 & 0 & 0 & 0 \\ 0 & 0 & B & -1 & 0 & 0 \\ 0 & 0 & 0 & B & -1 & 0 \\ 0 & 0 & 0 & 0 & B & -1 \\ 0 & 0 & 0 & 0 & 0 & B \end{pmatrix}, \quad (2.2)$$

and

$$B = 1 + a_5 \left(-\frac{a}{2} \nabla_\mu \nabla_\mu^* - \frac{m_0}{a} \right). \quad (2.3)$$

Due to its structure of the chiral hopping and the boundary condition in the flavor space, a single light Dirac fermion can emerge in the spectrum. The exact eigenvalues and eigenvectors of the mass matrix for the free theory at a finite flavor N has been given in [14]. This light fermion can be probed most suitably by the field variables at the boundary in the flavor space and is denoted as $q(x)$ and $\bar{q}(x)$ by Furman and Shamir:

$$q(x) = \psi_{1L}(x) + \psi_{NR}(x), \quad \bar{q}(x) = \bar{\psi}_{1L}(x) + \bar{\psi}_{NR}(x). \quad (2.4)$$

Following Neuberger, we may change the flavor index of the left-handed component by the chirally asymmetric parity transformation in the flavor space:

$$\psi'_s(x) = (P_R + P_L P)_{st} \psi_t(x), \quad (2.5)$$

$$\bar{\psi}'_s(x) = \bar{\psi}_t(x) (P_R P + P_L)_{ts}, \quad (2.6)$$

where

$$P_{st} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad (N = 6). \quad (2.7)$$

By this transformation, the mass matrix becomes hermitian,

$$M_{st}^H = M_{st} P = P M_{st}^\dagger$$

$$= \frac{1}{a_5} \begin{pmatrix} 0 & 0 & 0 & 0 & -1 & B \\ 0 & 0 & 0 & -1 & B & 0 \\ 0 & 0 & -1 & B & 0 & 0 \\ 0 & -1 & B & 0 & 0 & 0 \\ -1 & B & 0 & 0 & 0 & 0 \\ B & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad (N=6). \quad (2.8)$$

In this basis, it is the N -th flavor field that is most suitable to probe the light fermion:

$$q(x) = \psi'_N(x), \quad \bar{q}(x) = \bar{\psi}'_N(x). \quad (2.9)$$

The chiral transformation adopted by Shamir and Furman [13] is given in this hermitian basis as follows:

$$\delta\psi'_s(x) = (\Gamma_5)_{st} \psi'_t(x), \quad (2.10)$$

where Γ_5 is given (for $N=6$) by

$$(\Gamma_5)_{st} = \begin{pmatrix} -\gamma_5 & 0 & 0 & 0 & 0 & 0 \\ 0 & -\gamma_5 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\gamma_5 & 0 & 0 & 0 \\ 0 & 0 & 0 & \gamma_5 & 0 & 0 \\ 0 & 0 & 0 & 0 & \gamma_5 & 0 \\ 0 & 0 & 0 & 0 & 0 & \gamma_5 \end{pmatrix} \quad (N=6). \quad (2.11)$$

In particular, the light fermion transforms as Eq. (1.4).

$$\delta q(x) = \gamma_5 q(x), \quad \delta \bar{q}(x) = \bar{q}(x) \gamma_5.$$

With this definition of the chiral transformation, the chiral symmetry is broken only by the diagonal $\frac{N}{2}$ -th element of the hermitian mass matrix Eq. (2.8), i.e. the (diagonal) mass term of the $\frac{N}{2}$ -th flavor:

$$\{\Gamma_5 D'_{\text{DW}} + D'_{\text{DW}} \Gamma_5\}_{st} = \frac{2}{a_5} \gamma_5 \delta_{s\frac{N}{2}} \delta_{t\frac{N}{2}}. \quad (2.12)$$

2.2 Pauli-Villars field and truncated overlap Dirac operator

In order to take the limit of infinite flavors and to relate the domain-wall fermion to the Dirac fermion described by the overlap Dirac operator, it is suitable to introduce, as a Pauli-Villars field, the N -flavor Wilson-Dirac

bosons with the flavor-mixing mass matrix which is anti-periodic in the flavor space [23]:

$$M_{st}^{\text{PV}} = \frac{1}{a_5} \begin{pmatrix} 0 & 0 & 0 & 0 & -1 & B \\ 0 & 0 & 0 & -1 & B & 0 \\ 0 & 0 & -1 & B & 0 & 0 \\ 0 & -1 & B & 0 & 0 & 0 \\ -1 & B & 0 & 0 & 0 & 0 \\ B & 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \quad (2.13)$$

Then the total action of the domain-wall fermion with the subtraction of the Pauli-Villars field is given by

$$\begin{aligned} \bar{S}_{\text{DW}} &= \sum_{s,t=1}^N a^4 \sum_x \bar{\psi}'_s(x) \left\{ \gamma_\mu \frac{1}{2} (\nabla_\mu + \nabla_\mu^*) \delta_{st} + M_{st}^H \right\} \psi'_t(x) \\ &+ \sum_{s,t=1}^N a^4 \sum_x \bar{\phi}'_s(x) \left\{ \gamma_\mu \frac{1}{2} (\nabla_\mu + \nabla_\mu^*) \delta_{st} + M_{st}^{\text{PV}} \right\} \phi'_t(x). \end{aligned} \quad (2.14)$$

As shown by Neuberger in [14] that the partition function of this total system at a finite flavor N can be written by a single determinant of the truncated overlap Dirac operator:

$$\bar{Z}_{\text{DW}} = Z_{\text{DW}} Z_{\text{PV}} = \det aD_N. \quad (2.15)$$

The truncated overlap Dirac operator is defined by Eq. (1.5) through the transfer matrix, which is given explicitly in the chiral basis ⁵ as

$$T \equiv e^{-a_5 \bar{H}} = \begin{pmatrix} \frac{1}{B} & \frac{1}{B} C \\ -C^\dagger \frac{1}{B} & B + C^\dagger \frac{1}{B} C \end{pmatrix}, \quad (2.17)$$

where

$$C = a_5 \sigma_\mu \frac{1}{2} (\nabla_\mu + \nabla_\mu^*) \quad (2.18)$$

⁵ The gamma matrices in the chiral basis are chosen as follows in our convention,

$$\gamma_\mu = \begin{pmatrix} 0 & \sigma_\mu \\ \sigma_\mu^* & 0 \end{pmatrix}, \quad \gamma_5 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \sigma_\mu = (1, \sigma_1, \sigma_2, \sigma_3). \quad (2.16)$$

and B is given by Eq. (2.3). Thus the domain-wall fermion with the subtraction of the Pauli-Villars field is equivalent to the Dirac fermion described by the truncated overlap Dirac operator:

$$S_N = a^4 \sum_x \bar{\psi}(x) D_N \psi(x). \quad (2.19)$$

The overlap Dirac operator, which is obtained from D_N in the limit of the infinite flavors (but with a finite a_5), is denoted as $\tilde{D} \equiv \lim_{N \rightarrow \infty} D_N$. This reduces to D of Eq. (1.5) in the limit that a_5 vanishes.

2.3 Effective action of the light fermion of domain-wall fermion

Now we evaluate the effective action of the light fermion field by integrating out $N - 1$ heavy flavors of the domain-wall fermion:

$$S_N^{\text{eff}} = a^4 \sum_x \bar{q}(x) D_N^{\text{eff}} q(x). \quad (2.20)$$

As mentioned in the introduction, this can be achieved by calculating the propagator of $q(x)$, because it should be given by the inverse of the effective Dirac operator for the field variables $q(x)$ and $\bar{q}(x)$. In order to obtain the propagator of these fields, we introduce the sources for them,

$$a^4 \sum_x \{ \bar{J}(x) q(x) + \bar{q}(x) J(x) \}. \quad (2.21)$$

We first describe the case with four flavors ($N = 4$) for simplicity and then generalize the result to any flavors. In the chiral basis of the gamma matrices, the original action of the domain-wall fermion Eq. (2.1) can be written in the matrix form as

$$S_{\text{DW}} = a^4 \sum_x \bar{\Psi}(x) D_{\text{DW}} \Psi(x), \quad (N = 4) \quad (2.22)$$

where

$$\bar{\Psi}(x) = (\bar{\psi}_{1L} \quad \bar{\psi}_{1R} \quad \bar{\psi}_{2L} \quad \bar{\psi}_{2R} \quad \bar{\psi}_{3L} \quad \bar{\psi}_{3R} \quad \bar{\psi}_{4L} \quad \bar{\psi}_{4R}), \quad (2.23)$$

$$D_{\text{DW}} = \frac{1}{a_5} \begin{pmatrix} B & C & 0 & 0 & 0 & 0 & 0 & 0 \\ -C^\dagger & B & 0 & -1 & 0 & 0 & 0 & 0 \\ -1 & 0 & B & C & 0 & 0 & 0 & 0 \\ 0 & 0 & -C^\dagger & B & 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 & B & C & 0 & 0 \\ 0 & 0 & 0 & 0 & -C^\dagger & B & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 & 0 & B & C \\ 0 & 0 & 0 & 0 & 0 & 0 & -C^\dagger & B \end{pmatrix}, \quad (2.24)$$

$$\Psi(x) = \begin{pmatrix} \psi_{1R} \\ \psi_{1L} \\ \psi_{2R} \\ \psi_{2L} \\ \psi_{3R} \\ \psi_{3L} \\ \psi_{4R} \\ \psi_{4L} \end{pmatrix}. \quad (2.25)$$

Following Neuberger [14], we then make the Dirac operator almost upper triangle. We first exchange the right-handed component and the left-handed component of each flavor in $\Psi(x)$:

$$\begin{pmatrix} \psi_{tR} \\ \psi_{tL} \end{pmatrix} \Rightarrow \begin{pmatrix} \psi_{tL} \\ \psi_{tR} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \psi_{tR} \\ \psi_{tL} \end{pmatrix}, \quad (t = 1, 2, 3, 4). \quad (2.26)$$

Then the Dirac operator of the domain-wall fermion becomes

$$D_{\text{DW}} \Rightarrow \frac{1}{a_5} \begin{pmatrix} C & B & 0 & 0 & 0 & 0 & 0 & 0 \\ B & -C^\dagger & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & C & B & 0 & 0 & 0 & 0 \\ 0 & 0 & B & -C^\dagger & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & C & B & 0 & 0 \\ 0 & 0 & 0 & 0 & B & -C^\dagger & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & C & B \\ 0 & 0 & 0 & 0 & 0 & 0 & B & -C^\dagger \end{pmatrix}. \quad (2.27)$$

We further move the first row down to the last row:

$$\Rightarrow D''_{\text{DW}} \equiv \frac{1}{a_5} \begin{pmatrix} B & -C^\dagger & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & C & B & 0 & 0 & 0 & 0 \\ 0 & 0 & B & -C^\dagger & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & C & B & 0 & 0 \\ 0 & 0 & 0 & 0 & B & -C^\dagger & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & C & B \\ 0 & 0 & 0 & 0 & 0 & 0 & B & -C^\dagger \\ C & B & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \quad (2.28)$$

Accordingly, the components of the domain-wall fermion fields reads

$$\begin{aligned}\bar{\Psi}(x) \implies \bar{\Psi}''(x) &= \begin{pmatrix} \bar{\psi}_{1R}'' & \bar{\psi}_{1L}'' & \bar{\psi}_{2R}'' & \bar{\psi}_{2L}'' & \bar{\psi}_{3R}'' & \bar{\psi}_{3L}'' & \bar{\psi}_{4R}'' & \bar{\psi}_{4L}'' \end{pmatrix} \\ &\equiv \begin{pmatrix} \bar{\psi}_{1R} & \bar{\psi}_{2L} & \bar{\psi}_{2R} & \bar{\psi}_{3L} & \bar{\psi}_{3R} & \bar{\psi}_{4L} & \bar{\psi}_{4R} & \bar{\psi}_{1L} \end{pmatrix},\end{aligned}\tag{2.29}$$

$$\Psi(x) \implies \Psi''(x) = \begin{pmatrix} \psi_{1L}'' \\ \psi_{1R}'' \\ \psi_{2L}'' \\ \psi_{2R}'' \\ \psi_{3L}'' \\ \psi_{3R}'' \\ \psi_{4L}'' \\ \psi_{4R}'' \end{pmatrix} \equiv \begin{pmatrix} \psi_{1L} \\ \psi_{1R} \\ \psi_{2L} \\ \psi_{2R} \\ \psi_{3L} \\ \psi_{3R} \\ \psi_{4L} \\ \psi_{4R} \end{pmatrix}.\tag{2.30}$$

The sources for $q(x)$ and $\bar{q}(x)$ may be expressed in this upper-triangle basis as follows:

$$\begin{aligned}a^4 \sum_x \left\{ \bar{J}(x) \left[P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \psi_1''(x) + P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \psi_N''(x) \right] \right. \\ \left. + \bar{\psi}_N''(x) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} J(x) \right\}.\end{aligned}\tag{2.31}$$

In order to express the two by two blocks of D_{DW}'' , we introduce the following abbreviations

$$\alpha = \frac{1}{a_5} \begin{pmatrix} B & -C^\dagger \\ 0 & -1 \end{pmatrix}, \quad \beta = \frac{1}{a_5} \begin{pmatrix} -1 & 0 \\ C & B \end{pmatrix},\tag{2.32}$$

$$\alpha_0 = \frac{1}{a_5} \begin{pmatrix} B & -C^\dagger \\ 0 & 0 \end{pmatrix}, \quad \beta_0 = \frac{1}{a_5} \begin{pmatrix} 0 & 0 \\ C & B \end{pmatrix}.\tag{2.33}$$

In this basis it is now easy to integrate fields from the first flavor down to the last flavor.⁶ The terms which include the field variables of the first flavor are following (The summation over the lattice indices x with the measure factor a^4 is understood in the following equations.):

$$\left[\bar{J}(x) P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \bar{\psi}_N''(x) \beta_0 \right] \psi_1''(x)$$

⁶ Note that $q_L(x)$ is still in the first component of $\Psi''(x)$, although $\bar{q}_L(x)$ is in the N -th component of $\bar{\Psi}''(x)$. This is why we could not evaluate the effective action directly in this basis which is convenient for the integration.

$$\begin{aligned}
& +\bar{\psi}_1''(x) \alpha \psi_1''(x) \\
& +\bar{\psi}_1''(x) \beta \psi_2''(x).
\end{aligned} \tag{2.34}$$

After integrating the first flavor, the terms which include the second flavor are found as follows:

$$\begin{aligned}
& \left[\bar{J}(x) P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \bar{\psi}_N''(x) \beta_0 \right] (-\alpha^{-1} \beta) \psi_2''(x) \\
& +\bar{\psi}_2''(x) \alpha \psi_2''(x) \\
& +\bar{\psi}_2''(x) \beta \psi_3''(x).
\end{aligned} \tag{2.35}$$

The integration of the second flavor leaves the terms which include the third flavor as follows:

$$\begin{aligned}
& \left[\bar{J}(x) P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \bar{\psi}_N''(x) \beta_0 \right] (-\alpha^{-1} \beta)^2 \psi_3''(x) \\
& +\bar{\psi}_3''(x) \alpha \psi_3''(x) \\
& +\bar{\psi}_3''(x) \beta \psi_4''(x).
\end{aligned} \tag{2.36}$$

After the integration of the third flavor, only the forth flavor remains:

$$\begin{aligned}
& \bar{J}(x) \left[P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (-\alpha^{-1} \beta)^{N-1} + P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right] \psi_N''(x) \\
& +\bar{\psi}_N''(x) \left[\alpha_0 + \beta_0 (-\alpha^{-1} \beta)^{N-1} \right] \psi_N''(x) \\
& +\bar{\psi}_N''(x) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} J(x), \quad (N=4).
\end{aligned} \tag{2.37}$$

As easily seen, this result holds true for any flavors N .

Now noting

$$\begin{aligned}
\alpha_0 \alpha^{-1} &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \\
\beta_0 \beta^{-1} &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \\
(-\beta \alpha^{-1}) &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \gamma_5 e^{a_5 \tilde{H}} \gamma_5 \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},
\end{aligned} \tag{2.38}$$

we can evaluate the factors of the first and second terms of Eq. (2.37) as follows:

$$\left[P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (-\alpha^{-1} \beta)^{N-1} + P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right] \alpha^{-1}$$

$$= -a_5 \left[P_R + P_L e^{Na_5 \tilde{H}} \right] (-\gamma_5) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (2.39)$$

$$\begin{aligned} & \left[\alpha_0 + \beta_0 (-\alpha^{-1} \beta)^{N-1} \right] \alpha^{-1} \\ &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \left[P_L + P_R e^{Na_5 \tilde{H}} \right] (-\gamma_5) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \end{aligned} \quad (2.40)$$

From these results, we obtain

$$\begin{aligned} \langle q(x) \bar{q}(y) \rangle &= \frac{a_5}{a^4} \left(P_R + P_L e^{Na_5 \tilde{H}} \right) \frac{1}{P_L + P_R e^{Na_5 \tilde{H}}} \\ &= \frac{a_5}{a^4} \frac{1 - \gamma_5 \tanh a_5 \frac{N}{2} \tilde{H}}{1 + \gamma_5 \tanh a_5 \frac{N}{2} \tilde{H}}. \end{aligned} \quad (2.41)$$

Then, the effective action of the light fermion turns out to be given by the following effective Dirac operator:

$$\begin{aligned} D_N^{\text{eff}}(x, y) &= \frac{1}{a^4} \langle q(x) \bar{q}(y) \rangle^{-1} \\ &= \frac{1}{a_5} \frac{1 + \gamma_5 \tanh \frac{N}{2} a_5 \tilde{H}}{1 - \gamma_5 \tanh \frac{N}{2} a_5 \tilde{H}}. \end{aligned} \quad (2.42)$$

2.4 The effective action of the light fermion and the Ginsparg-Wilson relation

We may also consider the effective action of the N -th flavor of the bosonic Pauli-Villars field. As we see from Eq. (2.13), the difference between the mass matrices of the domain-wall fermion and the Pauli-Villars field is only in the diagonal N -th element of the latter, which takes account of the anti-periodicity in the flavor space. Then the integration of the first $N - 1$ components can be achieved just in the same way as the case of the domain-wall fermion. The effective action of the N -th flavor of the Pauli-Villars field turns out to be same as that of the light-fermion, but in this case, with the additional mass term. Thus the total effective action of the domain-wall fermion with the subtraction can be written in the following form:

$$\bar{S}_N^{\text{eff}} = a^4 \sum_x \bar{q}(x) D_N^{\text{eff}} q(x) + a^4 \sum_x \bar{Q}(x) \left\{ D_N^{\text{eff}} + \frac{1}{a_5} \right\} Q(x), \quad (2.43)$$

where we have denoted the N -th flavor of the Pauli-Villars field by

$$Q(x) = \phi'_N(x), \quad \bar{Q}(x) = \bar{\phi}'_N(x). \quad (2.44)$$

This result is consistent with Eq. (2.15), because the relation holds true

$$\frac{a_5 D_N^{\text{eff}}}{1 + a_5 D_N^{\text{eff}}} = a D_N, \quad (2.45)$$

and the partition function calculated from the effective action $\bar{S}_N^{\text{eff}}$ (Eq. (2.43)) is identical to that from the action with the truncated overlap Dirac operator S_N (Eq. (2.19)):

$$\begin{aligned} \bar{Z}_{\text{DW}} &= \int [dq d\bar{q}] [dQ d\bar{Q}] e^{-a^4 \sum_x \bar{q}(x) D_N^{\text{eff}} q(x) - a^4 \sum_x \bar{Q}(x) \left\{ D_N^{\text{eff}} + \frac{1}{a_5} \right\} Q(x)} \\ &= \int [d\psi d\bar{\psi}] e^{-a^4 \sum_x \bar{\psi}(x) D_N \psi(x)}. \end{aligned} \quad (2.46)$$

Eq. (2.45) may be written also in the following form:

$$\frac{1}{a_5} D_N^{\text{eff}-1} + \delta(x, y) = \frac{1}{a} D_N^{-1}. \quad (2.47)$$

Namely, the inverse of the effective Dirac operator (the propagator of the light fermion) gives the inverse of the truncated overlap Dirac operator up to a local contact term. This contact term just takes account of the chiral symmetry breaking in the Ginsparg-Wilson relation, which holds true for the overlap Dirac operator in the limit of the infinite flavors. This result implies that the propagator of the light fermion reduces to *the chirally symmetric part of the inverse of the overlap Dirac operator* in the limit of the infinite flavors, assuming that $\tilde{D}$ does not have any zero mode:

$$\gamma_5 D^{\text{eff}-1} + D^{\text{eff}-1} \gamma_5 = 0 \quad (2.48)$$

and

$$\tilde{D}^{-1} = \frac{a}{a_5} D^{\text{eff}-1} + a \delta(x, y) \quad (N = \infty). \quad (2.49)$$

It may be interesting to observe that the infinitely many flavors of the domain-wall fermion serve to prepare the chirally symmetric part of the inverse of the overlap Dirac operator and that it is the Pauli-Villars field (N -th flavor) that add it the contact term of the chiral symmetry breaking in the Ginsparg-Wilson relation. This contrasts with the situation of the original derivation of the Ginsparg-Wilson relation in [4], where the chiral symmetry breaking is introduced by the kernel of the block-spin transformation.

The properties of the chirally symmetric part of the inverse of the Dirac operator which satisfies the Ginsparg-Wilson relation have been discussed

extensively by Chiu et al. [24]. Eq. (2.45) reduces to the expression of the Dirac operator in terms of the chirally symmetric part discussed there.

As is also emphasized by these authors, the effective action of the light fermion, in the limit of infinite flavors, must necessarily be nonlocal. This can be easily checked in the free theory. Then, the effective action of the light fermion itself is not defined well and useful in this limit. In fact, a subtlety appears when one attempts to calculate the axial anomaly from the effective action, as will be discussed in the last section.

However, as we will discuss below, the light fermion field variables $q(x)$ and $\bar{q}(x)$ still remain to be a good probe for the Dirac fermion described by the overlap Dirac operator. They can be expressed by the Dirac fermion field $\psi(x)$ and $\bar{\psi}(x)$ through a certain local expression including the overlap Dirac operator $\bar{D}$. In this expression, we do not encounter any singularity associated with the nonlocal behavior of D^{eff} . Moreover, the low energy observables of the domain-wall fermion which are written in terms of $q(x)$ and $\bar{q}(x)$ can be expressed by $\psi(x)$ and $\bar{\psi}(x)$. And these observables, in fact, turn out to have good chiral properties from the point of view of the exact chiral symmetry based on the Ginsparg-Wilson relation.

3 Chiral property of the light fermion

With the result obtained in the previous section, we consider the relation between the light fermion field of the domain-wall fermion and the Dirac fermion field described by the truncated overlap Dirac operator. Then we discuss the chiral property of the light fermion field and various observables written by it in view of the exact chiral symmetry based on the Ginsparg-Wilson relation.

3.1 Light fermion field $q(x)$ and $\bar{q}(x)$

The correspondence between the light fermion field $q(x)$ and $\bar{q}(x)$ and the Dirac fermion field $\psi(x)$ and $\bar{\psi}(x)$ which is described by the truncated overlap Dirac operator is now easily understood. We may relate them by

$$\begin{aligned} q(x) &= Z \frac{1}{1 + a_5 D_N^{\text{eff}}} \psi(x) = Z \left(1 - \frac{a}{2} R D_N\right) \psi(x), \\ \bar{q}(x) &= \bar{\psi}(x), \end{aligned} \tag{3.1}$$

where $Z = \frac{a_5}{a}$ and $R = 2$. In the functional integral of the partition function, the Jacobian of the change of the field variable from $q(x)$ to $\psi(x)$ along this

relation, just compensates the determinant resulting from the integration of the last flavor of the bosonic Pauli-Villars field $Q(x)$.

A few comments are in order. As long as the equivalence of the partition functions is concerned, there is an ambiguity in the correspondence Eq. (3.1) which is related to the scale transformation:

$$\psi(x) \longrightarrow z \psi(x), \quad \bar{\psi}(x) \longrightarrow z^{-1} \bar{\psi}(x). \quad (3.2)$$

This scale factor may even depend on D_N^{eff} . We have fixed this freedom so that the chiral transformation of $\bar{q}(x)$ matches with that of $\bar{\psi}(x)$. As to the choice of the constant scale factor, it is our convention for simplicity. The correspondence Eq. (3.1) would be different if we adopt a different transformation of the exact chiral symmetry of Lüscher from Eq. (1.2).

3.2 Chiral transformation of the light fermion

The explicit relation between $q(x)$ and $\psi(x)$ helps us to understand the chiral properties of the domain-wall fermion in terms of the (would-be) exact chiral symmetry of the Ginsparg-Wilson fermion described by the (truncated) overlap Dirac operator. First of all, Eq. (2.47) relates quantitatively the chiral symmetry breaking in the domain-wall fermion to the breaking of the Ginsparg-Wilson relation in the truncated overlap Dirac fermion: these breakings can be characterized by a single quantity

$$\delta_N \equiv Z^{-1} \left(\gamma_5 D_N^{\text{eff}-1} + D_N^{\text{eff}-1} \gamma_5 \right) \quad (3.3)$$

$$\equiv \gamma_5 D_N^{-1} + D_N^{-1} \gamma_5 - aR\gamma_5. \quad (3.4)$$

We may consider the chiral transformation for the Dirac fermion described by the truncated overlap Dirac operator à la Lüscher

$$\delta\psi(x) = \gamma_5 (1 - aR D_N) \psi(x), \quad \delta\bar{\psi}(x) = \bar{\psi}(x) \gamma_5. \quad (3.5)$$

The chiral symmetry of the action is broken by the amount

$$\delta S_N = a^4 \sum_x \bar{\psi}(x) \Delta_N \psi(x), \quad (3.6)$$

$$\Delta_N \equiv \gamma_5 \left\{ 1 - \left(\tanh \frac{N}{2} a_5 \tilde{H} \right)^2 \right\} = D_N \delta_N D_N. \quad (3.7)$$

This breaking vanishes in the limit of infinite flavors $N = \infty$, as long as the eigenvalues of $\tilde{H}$ are bounded from zero uniformly with respect to the gauge

fields in consideration. We assume this in the following discussions. We also assume that $\tilde{D}$, the overlap Dirac operator in the limit of the infinite flavors, should not have any zero mode and that its inverse should exist, in order to assure that δ_N should vanish in this limit, too. When we discuss the anomaly of the domain-wall fermion in relation to the index of $\tilde{D}$, we introduce the mass term of the light fermion to assure the limit.

The chiral transformation of Eq. (3.5) induces the transformation of the light fermion as follows:

$$\begin{aligned}
\delta q(x) &= Z \left(1 - \frac{a}{2} R D_N\right) \delta \psi(x) \\
&= Z \left(1 - \frac{a}{2} R D_N\right) \gamma_5 (1 - a R D_N) \psi(x) \\
&= Z \left\{ \gamma_5 (1 - a R D_N) - \frac{a}{2} R D_N \gamma_5 (1 - a R D_N) \right\} \psi(x) \\
&= Z \left\{ \gamma_5 \left(1 - \frac{a}{2} R D_N\right) - \frac{a}{2} R \Delta_N \right\} \psi(x) \\
&= \gamma_5 q(x) - a_5 \Delta_N \psi(x).
\end{aligned} \tag{3.8}$$

And it reduces to the transformation of Eq. (1.4) in the limit of the infinite flavors. Thus we can see how the chiral transformation of Eq. (1.4), which is adopted by Furman and Shamir (See also Eq. (2.10)), is related to that of Lüscher Eq. (1.2) in the limit of the infinite flavors.

3.3 Low energy observables in terms of the light fermion

Next we consider the low energy observables of the domain-wall fermion, which are written in terms of $q(x)$ and $\bar{q}(x)$. We discuss the relation to the observables in terms of the Dirac fermion field which is described by the overlap Dirac operator and whose chiral property is governed by the Ginsparg-Wilson relation.

3.3.1 Scalar and pseudo scalar bilinear operators

First of all, from the explicit relation Eq. (3.1) between the light fermion field and the Dirac field described by the truncated overlap Dirac operator, we obtain the relation of the scalar and pseudo scalar bilinear operators:

$$\bar{q}(x)q(x) = Z \bar{\psi}(x) \left(1 - \frac{a}{2} R D_N\right) \psi(x), \tag{3.9}$$

$$\bar{q}(x)\gamma_5 q(x) = Z \bar{\psi}(x)\gamma_5 \left(1 - \frac{a}{2} R D_N\right) \psi(x). \tag{3.10}$$

In this respect, it is interesting to note that the scalar and pseudo scalar bilinear operators in the r.h.s. consist the exact chiral multiplet of the chiral transformation Eq. (1.2) in the limit of infinite flavors, as discussed by Niedermayer [25]. In fact, these operators can be written in the chiral components defined by $\hat{\gamma}_5 = \gamma_5 \left(1 - aR\tilde{D}\right)$ for $\psi(x)$ and by γ_5 for $\bar{\psi}(x)$ as follows:

$$\bar{q}(x)q(x) \longrightarrow \bar{\psi}_L(x)\psi_R(x) + \bar{\psi}_R(x)\psi_L(x), \quad (3.11)$$

$$\bar{q}(x)\gamma_5 q(x) \longrightarrow \bar{\psi}_L(x)\psi_R(x) - \bar{\psi}_R(x)\psi_L(x). \quad (3.12)$$

3.3.2 Conserved vector currents

The vector current of the domain-wall fermion is conserved at a finite flavor N . It is expected to correspond to the conserved vector current of the truncated overlap Dirac fermion. We will show that the vector current of the domain-wall fermion, if it is probed by the light fermion field $q(x)$ and $\bar{q}(x)$ at low energy, just corresponds to the conserved vector current of the truncated overlap Dirac fermion.

In general, a vector current may be defined with the kernel

$$V_\mu^a(x) = \sum_{y,z} \bar{\psi}(x) K_\mu^a(x; y, z) \psi(x). \quad (3.13)$$

The kernel is obtained from the Dirac operator by introducing the auxiliary vector field

$$U_\mu^B(x) = \exp \left(i B_\mu^a(x) T^a \right), \quad (3.14)$$

where T^a are assumed as the generators of the $U(N_F)$ real flavor group, and by differentiating the Dirac operator with respect to the vector field:

$$\delta D(y, z) = \sum_x B_\mu^a(x) K_\mu^a(x; y, z) + \mathcal{O}(B^2). \quad (3.15)$$

By this procedure, we can obtain the kernel for the domain-wall fermion from $D'_{\text{DW}} \equiv \gamma_\mu \frac{1}{2} (\nabla_\mu + \nabla_\mu^*) + M^H$, which we denote as $K_{\mu\text{DW}}^a$. The kernel for the truncated overlap Dirac fermion is obtained from D_N , which we denote as $K_{\mu N}^a$. Then, the vector current of the domain-wall fermion with the subtraction can be written as

$$V_{\mu\text{DW}}^a(x) = \sum_{st}^N \bar{\psi}'_s \{ K_{\mu\text{DW}}^a(x) \}_{st} \psi'_t. \quad (3.16)$$

The vector current of the truncated overlap Dirac fermion can be written as

$$V_N^a(x) = \bar{\psi} K_{\mu N}^a(x) \psi. \quad (3.17)$$

For these two vector currents, we can infer the following identity:

$$\langle q(y) V_{\mu \text{DW}}^a(x) \bar{q}(z) \rangle_c = Z \langle \psi(y) V_{\mu N}^a(x) \bar{\psi}(z) \rangle_c. \quad (3.18)$$

This identity follows from the relation between D_{DW}^{-1} and D_N^{-1} given by Eq. (2.47): in fact, it follows that

$$\left\{ \delta D_{\text{DW}}'^{-1} \right\}_{NN} = \delta D_N^{\text{eff}^{-1}} = Z \delta D_N^{-1}. \quad (3.19)$$

Using the identity $\delta D^{-1} = -D^{-1} \delta D D^{-1}$, it can be expressed as Eq. (3.18).

3.3.3 Almost conserved axial vector currents

According to Shamir and Furman [13], the axial vector current of the domain-wall fermion is conserved up to the corrections suppressed exponentially in the number of flavors. Then it is expected that this axial vector current is related in the limit of the infinite flavors to the conserved axial vector current of the overlap Dirac fermion. We will show that the axial vector current of the domain-wall fermion, if it is probed by the light fermion field $q(x)$ and $\bar{q}(x)$ at low energy, reduces to the conserved axial vector current of the overlap Dirac fermion.

The axial vector current of the domain-wall fermion is defined with the kernel of the vector current as

$$A_{\mu \text{DW}}^a(x) = \sum_{s,t}^N \bar{\psi}_s' \{ K_{\mu \text{DW}}^a(x) \Gamma_5 \}_{st} \psi_t'. \quad (3.20)$$

On the other hand, the axial vector current of the truncated overlap Dirac fermion may be defined as

$$A_{\mu N}^a(x) = \bar{\psi} K_{\mu N}^a(x) \gamma_5 (1 - a R D_N) \psi, \quad (3.21)$$

which naturally follows from the chiral transformation Eq. (3.5). This current reduces to the Noether current associated with the exact chiral symmetry of Lüscher [29] in the limit of the infinite flavors, which we denote as

$$\tilde{A}_\mu^a(x) = \bar{\psi} \tilde{K}_\mu^a(x) \gamma_5 (1 - a R \tilde{D}) \psi. \quad (3.22)$$

For these two axial vector currents, we can infer the following relation:

$$\lim_{N \rightarrow \infty} \langle q(y) A_{\mu\text{DW}}^a(x) \bar{q}(z) \rangle_c = Z \langle \psi(y) \tilde{A}_\mu^a(x) \bar{\psi}(z) \rangle_c. \quad (3.23)$$

This relation can be shown in the following way. The Dirac operator of the domain-wall fermion satisfies

$$\{\Gamma_5 D'_{\text{DW}} + D'_{\text{DW}} \Gamma_5\}_{st} = 2\gamma_5 \delta_{s\frac{N}{2}} \delta_{t\frac{N}{2}}. \quad (3.24)$$

On the other hand, the truncated overlap Dirac operator satisfies

$$\gamma_5 D_N + D_N \gamma_5 (1 - aR D_N) = \Delta_N. \quad (3.25)$$

Then it turns out that the vector and axial vector currents are related each other as follows:

$$\begin{aligned} & \langle q(y) A_{\mu\text{DW}}^a(x) \bar{q}(z) \rangle + \langle q(y) V_{\mu\text{DW}}^a(x) \bar{q}(z) \rangle \gamma_5 \\ &= \sum_s^N \{D_{\text{DW}}^{-1} K_{\mu\text{DW}}^a(x)\}_{N,s} \cdot \{D_{\text{DW}}^{-1}\}_{s,\frac{N}{2}} 2\gamma_5 \{D_{\text{DW}}^{-1}\}_{\frac{N}{2},N}(y,z), \end{aligned} \quad (3.26)$$

and

$$\begin{aligned} & \langle \psi(y) A_{\mu N}^a(x) \bar{\psi}(z) \rangle + \langle \psi(y) V_{\mu N}^a(x) \bar{\psi}(z) \rangle \gamma_5 \\ &= \{D_N^{-1} K_{\mu N}(x)\} D_N^{-1} \Delta_N D_N^{-1}(y,z). \end{aligned} \quad (3.27)$$

Using Eq. (3.18), we obtain

$$\begin{aligned} & \langle q(y) A_{\mu\text{DW}}^a(x) \bar{q}(z) \rangle \\ &= Z \langle \psi(y) A_{\mu N}^a(x) \bar{\psi}(z) \rangle - Z D_N^{-1} K_{\mu N}(x) D_N^{-1} \Delta_N D_N^{-1}(y,z) \\ & \quad + \{D_{\text{DW}}^{-1} K_{\mu\text{DW}}(x) D_{\text{DW}}^{-1}\}_{N,\frac{N}{2}} 2\gamma_5 \{D_{\text{DW}}^{-1}\}_{\frac{N}{2},N}(y,z). \end{aligned} \quad (3.28)$$

Now, it is possible to evaluate the propagator between the $\frac{N}{2}$ -th flavor and the N -th flavor of the domain-wall fermion, using the similar method to evaluate the propagator of the light fermion in section 2. The calculation is described in the appendix A. The result can be expressed as

$$\begin{aligned} \{D_{\text{DW}}\}_{\frac{N}{2},N}^{-1}(x,y) &\equiv \langle \psi'_{\frac{N}{2}}(x) \bar{q}(y) \rangle \\ &= \frac{1}{2 \cosh \frac{N}{2} a_5 \tilde{H}} D_N^{-1}(x,y). \end{aligned} \quad (3.29)$$

From this and Eq. (3.7), we infer that the second and third terms of the right-hand sides of Eq. (3.28) vanish in the limit of the infinite flavors and we finally obtain Eq. (3.23).

Thus the vacuum expectation value of the axial vector current of the domain-wall fermion with respect to the light fermion field leads directly to that of the axial vector current associated with the exact chiral symmetry based on the Ginsparg-Wilson relation. Note that our results hold true with dynamical gauge fields as long as the eigenvalues of $\tilde{H}$ are bounded from zero uniformly with respect to the gauge fields. The renormalization factor of the axial current of the domain-wall fermion thus reduces to unity in the limit of the infinite flavors. Our result is consistent with the perturbative result at one-loop obtained by Aoki and Taniguchi [20].

3.3.4 Correspondence of various observables

From the above results, we can obtain the correspondence of the various observables of the light fermion of the domain-wall fermion to the observables of the Dirac fermion which is described by the (truncated) overlap Dirac operator. We summarize it here. (Note again that in this paper, the bracket symbol $\langle \cdots \rangle$ denotes the Wick contraction of the fermion fields in it by their propagators, not the fermionic vacuum expectation values which must includes the weight of the fermion action. The bra-ket symbol with the subscript c denotes the connected contraction.)

- Correlation of the vector currents:

$$\left\langle V_{\mu\text{DW}}^a(x) \bar{q}(y) \gamma_\nu T^b q(y) \right\rangle_c = Z \left\langle V_{\mu N}^a(x) \bar{\psi}(y) \gamma_\nu T^b \psi(y) \right\rangle_c. \quad (3.30)$$

- Correlation of the pseudo scalar densities:

$$\begin{aligned} & \left\langle \bar{q}(x) \gamma_5 T^a q(x) \bar{q}(y) \gamma_5 T^b q(y) \right\rangle \\ &= Z^2 \left\langle \bar{\psi}(x) \gamma_5 \left(1 - \frac{a}{2} R D_N\right) T^a \psi(x) \bar{\psi}(y) \gamma_5 \left(1 - \frac{a}{2} R D_N\right) T^b \psi(y) \right\rangle. \end{aligned} \quad (3.31)$$

- Correlation of the axial vector current and pseudo scalar density:

$$\lim_{N \rightarrow \infty} \left\langle A_{\mu\text{DW}}^a(x) \bar{q}(y) \gamma_5 T^b q(y) \right\rangle_c = Z \left\langle \tilde{A}_\mu^a(x) \bar{\psi}(y) \gamma_5 T^b \psi(y) \right\rangle_c. \quad (3.32)$$

- Amplitude of K^0 - $\bar{K}^0$ mixing:

$$\begin{aligned} \lim_{N \rightarrow \infty} \left\langle \bar{d}_N s_N(y) \left(\sum_{st}^N \bar{s}_s \{K_{\mu\text{DW}}(x) (1 - \Gamma_5)\}_{st} d_t \right)^2 \bar{d}_N s_N(z) \right\rangle \\ = Z^2 \left\langle \bar{d}s(y) \left(\bar{s} \tilde{K}_\mu(x) (1 - \hat{\gamma}_5) d \right)^2 \bar{d}s(z) \right\rangle, \end{aligned} \quad (3.33)$$

where $s_t(x)$ and $d_t(x)$ ($t = 1, \dots, N$) are the domain-wall fermion fields for s-quark and d-quark, respectively. $s(x)$ and $d(x)$ are the Dirac fermion fields which are described by the overlap Dirac operator $\tilde{D}$ for s-quark and d-quark, respectively.

Thus we see that the observables in the light fermion field variables $q(x)$ and $\bar{q}(x)$ of the domain-wall fermion (with the subtraction of the Pauli-Villars field) leads directly to the observables which have good chiral property with respect to the exact chiral symmetry based on the Ginsparg-Wilson relation.

4 Axial anomaly of domain-wall fermion

The explicit chiral symmetry breaking in the domain-wall fermion under the chiral transformation Eq. (2.10) is expected to reproduce the axial anomaly [26]. Here we examine this chiral symmetry breaking term in relation to the axial anomaly which is associated with the exact chiral symmetry based on the Ginsparg-Wilson relation [27, 5],

$$- a \text{tr} \gamma_5 R \tilde{D}(x, x). \quad (4.1)$$

4.1 Axial anomaly

With the definition of the chiral transformation Eq. (2.10), the chiral symmetry breaking occurs at the diagonal $\frac{N}{2}$ -th element of the hermitian mass matrix Eq. (2.8), i.e. mass term of the $\frac{N}{2}$ -th flavor. In fact, the flavor singlet axial vector current of the domain-wall fermion satisfies the axial Ward-Takahashi identity:

$$\partial_\mu^* \langle A_{\mu\text{DW}}(x) \rangle = \frac{2}{a_5} \left\langle \bar{\psi}'_{\frac{N}{2}}(x) \gamma_5 \psi'_{\frac{N}{2}}(x) \right\rangle. \quad (4.2)$$

It is expected that the explicit breaking in the r.h.s. of the above identity should reproduce the axial anomaly. It has been shown by a perturbative calculation that it is indeed the case [26].

It is actually possible to evaluate the breaking term non-perturbatively. Using the similar method to evaluate the propagator of the light fermion in section 2, the propagator of the $\frac{N}{2}$ -th flavor fermion of the domain-wall fermion is evaluated explicitly as

$$\begin{aligned} \left\langle \psi'_{\frac{N}{2}}(x) \bar{\psi}'_{\frac{N}{2}}(y) \right\rangle &= \frac{a_5}{a^4} \left(-1 + \gamma_5 \frac{1}{2} a R D_N \gamma_5 \right. \\ &\quad \left. + \frac{1}{2a} \frac{1}{\cosh \frac{N}{2} a_5 \tilde{H}} D_N^{-1} \gamma_5 \frac{1}{\cosh \frac{N}{2} a_5 \tilde{H}} \gamma_5 \right). \end{aligned} \quad (4.3)$$

The calculation is described in the appendix A. Then it follows immediately that the chiral symmetry breaking term can be written as

$$\begin{aligned} \frac{2}{a_5} \left\langle \bar{\psi}'_{\frac{N}{2}}(x) \gamma_5 \psi'_{\frac{N}{2}}(x) \right\rangle &= \frac{1}{a^4} \left(-a \text{tr} \gamma_5 R D_N(x, x) \right. \\ &\quad \left. - \text{tr} D_N^{-1} \gamma_5 \frac{1}{a \cosh^2 \frac{N}{2} a_5 \tilde{H}} (x, x) \right) \\ &= \frac{1}{a^4} \left(-a \text{tr} \gamma_5 R D_N(x, x) - \text{tr} D_N^{-1} \Delta_N(x, x) \right). \end{aligned} \quad (4.4)$$

We can see that it reduces to the anomaly Eq. (4.1) in the limit of the infinite flavors.

4.2 Relation to the index of overlap Dirac operator

We next discuss the relation of the axial anomaly of the domain-wall fermion to the index of the overlap Dirac operator $\tilde{D}$. For this purpose, we consider the situation so that $\tilde{D}$ has zero modes. In order to make the limit of the infinite flavors well-defined, we introduce the mass term of the light fermion.

$$m \bar{q}(x) q(x). \quad (4.5)$$

Then it follows from the axial Ward-Takahashi identity that

$$a^4 \sum_x 2 \left\langle \bar{\psi}'_{\frac{N}{2}}(x) \gamma_5 \psi'_{\frac{N}{2}}(x) \right\rangle + a^4 \sum_x 2m \langle \bar{q}(x) \gamma_5 q(x) \rangle = 0. \quad (4.6)$$

The second term of the l.h.s. is evaluated as

$$\begin{aligned}
-a^4 \sum_x 2m \langle \bar{q}(x) \gamma_5 q(x) \rangle &= 2m \text{Tr} \gamma_5 \frac{1}{D_N^{\text{eff}} + m} \\
&= 2m \text{Tr} \gamma_5 \left(1 - \frac{a}{2} R D_N \right) \frac{1}{D_N + m \left(1 - \frac{a}{2} R D_N \right)} \\
&= -a \text{Tr} \gamma_5 R D_N \\
&\quad - (1 - am) \text{Tr} \Delta_N \frac{1}{D_N + m \left(1 - \frac{a}{2} R D_N \right)}. \tag{4.7}
\end{aligned}$$

Therefore we obtain in the limit of the infinite flavors

$$\lim_{N \rightarrow \infty} a^4 \sum_x 2 \left\langle \bar{\psi}'_{\frac{N}{2}}(x) \gamma_5 \psi'_{\frac{N}{2}}(x) \right\rangle = -a \text{Tr} \gamma_5 R \tilde{D} = 2 \text{Index} \left(\tilde{D} \right). \tag{4.8}$$

Since all the dependence on m comes with Δ_N , this limit does not depend on the value of m .

This explicit chiral symmetry breaking term has been evaluated numerically by Argonne group [19] as the probe of the topological charge. We have seen how this quantity is related to the index of the overlap Dirac operator which satisfies the Ginsparg-Wilson relation.

5 Contributions of the Pauli-Villars field

Finally, we will discuss the contributions of the Pauli-Villars field to vector and axial vector currents and axial anomaly. In the previous sections, we have discussed the currents of the domain-wall fermion probed by the light fermion field $q(x)$ and $\bar{q}(x)$. However, if we consider the currents themselves and their correspondence to those of the overlap Dirac fermion, we need to include the contribution of the Pauli-Villars fields.

5.1 Vector and axial vector currents

The conserved vector current of the domain-wall fermion with the subtraction of the Pauli-Villars fields is defined by

$$\bar{V}_{\mu\text{DW}}^a(x) = \sum_{st}^N \bar{\psi}'_s \{ K_{\mu\text{DW}}^a(x) \}_{st} \psi'_t + \sum_{st}^N \bar{\phi}'_s \{ K_{\mu\text{DW}}^a(x) \}_{st} \phi'_t. \tag{5.1}$$

Note that we obtain the same kernel for the Pauli-Villars field as the domain-wall fermion, from $D'_{\text{PV}} \equiv \gamma_\mu \frac{1}{2} (\nabla_\mu + \nabla_\mu^*) + M^{\text{PV}}$. The almost conserved axial vector current is defined similarly by

$$\bar{A}_{\mu\text{DW}}^a(x) = \sum_{s,t}^N \bar{\psi}'_s \{K_{\mu\text{DW}}^a(x) \Gamma_5\}_{st} \psi'_t + \sum_{s,t}^N \bar{\phi}'_s \{K_{\mu\text{DW}}^a(x) \Gamma_5\}_{st} \phi'_t. \quad (5.2)$$

For these vector and axial vector currents, we can infer the following identities:

$$\langle \bar{V}_{\mu\text{DW}}^a(x) \rangle = \langle V_{\mu N}^a(x) \rangle, \quad (5.3)$$

$$\lim_{N \rightarrow \infty} \langle \bar{A}_{\mu\text{DW}}^a(x) \rangle = \langle \tilde{A}_\mu^a(x) \rangle. \quad (5.4)$$

The identity of the vector currents follows immediately from Eq. (2.15). But we will show it here directly for later use. The l.h.s. of Eq. (5.3), if multiplied with the auxiliary vector field $B_\mu^a(x)$, can be rewritten as follows:

$$\begin{aligned} (\text{l.h.s.}) &\simeq a^4 \sum_x B_\mu^a(x) \langle \bar{V}_{\mu\text{DW}}^a(x) \rangle \\ &= - \sum_t^N \text{tr} \left\{ \delta D'_{\text{DW}} \left(D'_{\text{DW}}{}^{-1} - D'_{\text{PV}}{}^{-1} \right) \right\}_{tt} \\ &= - \sum_t^N \text{tr} \left\{ \delta D'_{\text{DW}} D'_{\text{DW}}{}^{-1} (M^{\text{PV}} - M^{\text{H}}) D'_{\text{PV}}{}^{-1} \right\}_{tt} \\ &= \sum_t^N \text{tr} \left\{ [D'_{\text{PV}} - (M^{\text{PV}} - M^{\text{H}})] \delta D'_{\text{DW}}{}^{-1} (M^{\text{PV}} - M^{\text{H}}) D'_{\text{PV}}{}^{-1} \right\}_{tt} \\ &= \text{tr} \left\{ \delta D'_{\text{DW}}{}^{-1} \right\}_{NN} \frac{1}{a_5} \left\{ 1 - \frac{1}{a_5} D'_{\text{PV}}{}^{-1} \right\}_{NN}. \end{aligned} \quad (5.5)$$

In the last equality, we have noted that

$$\{M^{\text{PV}} - M^{\text{H}}\}_{st} = \frac{1}{a_5} \delta_{sN} \delta_{Nt}. \quad (5.6)$$

The propagator of the N -th flavor of the Pauli-Villars field has been evaluated in section 2 as

$$\frac{1}{a_5} \left\{ D'_{\text{PV}}{}^{-1} \right\}_{NN} = \frac{a^4}{a_5} \langle Q(x) \bar{Q}(y) \rangle = \frac{1}{a_5 D_N^{\text{eff}} + 1} = (1 - a D_N). \quad (5.7)$$

With this result and Eq. (3.19), we obtain

$$\begin{aligned}
(\text{l.h.s.}) &= \text{tr} \delta D_N^{-1} D_N \\
&= a^4 \sum_x B_\mu^a(x) \langle V_{\mu N}^a(x) \rangle.
\end{aligned} \tag{5.8}$$

As to the identity of the axial vector current, using Eqs. (2.12) and (3.25) and through the similar calculation as in Eq. (5.5), we obtain

$$\begin{aligned}
\langle \bar{A}_{\mu \text{DW}}^a(x) \rangle &= \langle A_{\mu N}^a(x) \rangle + \text{tr} K_{\mu N}^a(x) D_N^{-1} \Delta_N D_N^{-1} \\
&\quad - \text{tr} \{ D_{\text{DW}}^{-1} K_{\mu \text{DW}}^a(x) D_{\text{DW}}^{-1} \}_{N, \frac{N}{2}} 2\gamma_5 \{ D_{\text{DW}}^{-1} \}_{\frac{N}{2}, N} D_N.
\end{aligned} \tag{5.9}$$

From this and Eq. (3.7), we infer that the second and third terms of the right-hand side of Eq. (5.9) vanish in the limit of the infinite flavors and we finally obtain Eq. (5.4).

5.2 Axial anomaly

The flavor singlet axial vector current of the domain-wall fermion with the subtraction of the Pauli-Villars field satisfies the following axial Ward-Takahashi identity:

$$\begin{aligned}
\partial_\mu^* \langle \bar{A}_{\mu \text{DW}}(x) \rangle &= 2 \left\langle \bar{\psi}'_{\frac{N}{2}}(x) \gamma_5 \psi'_{\frac{N}{2}}(x) \right\rangle + 2 \left\langle \bar{\phi}'_{\frac{N}{2}}(x) \gamma_5 \phi'_{\frac{N}{2}}(x) \right\rangle + 2 \langle \bar{Q}(x) \gamma_5 Q(x) \rangle,
\end{aligned} \tag{5.10}$$

where $Q(x)$ and $\bar{Q}(x)$ stand for the N -th flavor of the Pauli-Villars field as defined in Eq. (2.44). Note that in this case three chiral symmetry breaking terms would contribute to axial anomaly. The first term is the contribution of the domain-wall fermion and was evaluated in section 4. The second and third terms are the contributions of the Pauli-Villars field. In order to evaluate them, we need the propagators of the $\frac{N}{2}$ -th flavor and N -th flavor of the Pauli-Villars field. The propagator of $Q(x)$ and $\bar{Q}(x)$ was evaluated in section 2.

$$\langle Q(x) \bar{Q}(y) \rangle = \frac{1}{a^4} \frac{1}{D_N^{\text{eff}} + \frac{1}{a_5}} = \frac{a_5}{a^4} (1 - a D_N). \tag{5.11}$$

The propagator of the $\frac{N}{2}$ -th flavor of the Pauli-Villars field can be evaluated using the similar method to evaluate the propagator of the $\frac{N}{2}$ -th flavor of the domain-wall fermion, which was used in section 4. The calculation is described in appendix A. The result can be written as

$$\left\langle \phi'_{\frac{N}{2}}(x) \bar{\phi}'_{\frac{N}{2}}(y) \right\rangle = -\gamma_5 \frac{a_5}{a^4} (1 - aD_N) \gamma_5. \quad (5.12)$$

From these results, we see that the net contribution of the anomalous terms of the r.h.s. of Eq. (5.10) is same as that of the domain-wall fermion (without the subtraction of the Pauli-Villars field) and is given by

$$\partial_\mu^* \langle \bar{A}_{\mu\text{DW}}(x) \rangle = \frac{1}{a^4} \left(-a \text{tr} \gamma_5 R D_N(x, x) - \text{tr} D_N^{-1} \Delta_N(x, x) \right). \quad (5.13)$$

This identity corresponds term by term to the axial Ward-Takahashi identity of the truncated overlap Dirac fermion, which is derived under the chiral transformation Eq. (3.5):

$$\partial_\mu^* \langle A_{\mu N}(x) \rangle = \frac{1}{a^4} \left(-a \text{tr} \gamma_5 R D_N(x, x) + \langle \bar{\psi}(x) \Delta_N \psi(x) \rangle \right). \quad (5.14)$$

And it reduces in the limit of the infinite flavors to the anomalous axial Ward-Takahashi identity of the overlap Dirac fermion:

$$\partial_\mu^* \langle \tilde{A}_\mu(x) \rangle = \frac{1}{a^4} \left(-a \text{tr} \gamma_5 R \tilde{D}(x, x) \right). \quad (5.15)$$

5.3 Probe for topological charge at a finite flavor

In view of the Eqs. (4.6), (4.7) and (4.8), it seems reasonable to probe the topological charge by the anomalous term of Eq. (4.2), as in the numerical calculation by Argonne group [19]. In this respect, we may probe the topological charge rather directly using the truncated overlap Dirac operator.

$$Q_N = -\frac{1}{2} a \text{Tr} \gamma_5 R D_N. \quad (5.16)$$

The topological charge density, $-\frac{1}{2} a \text{tr} \gamma_5 R D_N(x, x)$ is assumed to be a local functional of the gauge fields in the limit of the infinite flavors [27, 5, 3]. This fact is reflected in the domain-wall fermion in that Q_N can be evaluated as a contribution of the massive flavor to the anomalous term of Eq. (4.2). We point out that Q_N can also be expressed as follows:

$$\begin{aligned} Q_N &= -a^4 \sum_x m \langle \bar{q}(x) \gamma_5 q(x) \rangle \Big|_{ma=1} \\ &= a^4 \sum_x \langle \bar{Q}(x) \gamma_5 Q(x) \rangle. \end{aligned} \quad (5.17)$$

6 Discussion

We have discussed the chiral property of the light fermion of the domain-wall fermion by considering its low energy effective action

$$S_N^{\text{eff}} = a^4 \sum_x \bar{q}(x) D_N^{\text{eff}} q(x), \quad (6.18)$$

where the effective Dirac operator has a simple relation to the truncated overlap Dirac operator as

$$\frac{a}{a_5} D_N^{\text{eff}^{-1}} + a\delta(x, y) = D_N^{-1}. \quad (6.19)$$

We have argued that the chiral property of the light fermion field is understandable also from the point of view of the exact chiral symmetry based on the Ginsparg-Wilson relation, which holds true for the overlap Dirac operator.

As discussed in section 2, in the limit of the infinite flavors, the effective action itself becomes chiral, but non-local. A subtlety of the effective action in this limit becomes clear when one attempts to calculate the axial anomaly directly from the effective action: with such chiral and non-local action, it is not easy to identify the source of the axial anomaly. One possible way might be to consider the explicit breaking term at a finite flavor N :

$$a^4 \sum_x \alpha(x) \left\langle \bar{q}(x) \left\{ \gamma_5, D_N^{\text{eff}} \right\} q(x) \right\rangle \Big|_{ma}. \quad (6.20)$$

(We have introduced a bare mass of the light fermion and $\alpha(x)$ is an infinitesimal local parameter.) In view of Eq. (4.7) and the axial Ward-Takahashi identity, as least for the global case this term actually reproduces the anomaly as the index of the overlap Dirac operator in the limit of the infinite flavors. We leave this question for future study.

Acknowledgments

We would like to thank A. Yamada, T. Onogi, S. Aoki, T. Izubuchi and Y. Taniguchi for enlightening discussions. Y.K. is also grateful to O. Miyamura and A. Nakamura for discussions. Y.K. is supported in part by Grant-in-Aid for Scientific Research from Ministry of Education, Science and Culture (#10740116, #10140214).

A Evaluation of propagators of heavy fermions and bosons

In this appendix, we evaluate the correlation function between the N -th flavor and the $\frac{N}{2}$ -th flavor of the domain-wall fermion

$$\left\langle \psi'_{\frac{N}{2}}(x) \bar{q}(y) \right\rangle, \quad (\text{A.21})$$

by the method used in section 2.3. This correlation function is used in the evaluation of the vacuum expectation value of the axial current in section 3. We also evaluate the correlation function of the $\frac{N}{2}$ -th flavor of the domain-wall fermion and that of the Pauli-Villars field

$$\left\langle \psi'_{\frac{N}{2}}(x) \bar{\psi}'_{\frac{N}{2}}(y) \right\rangle, \quad (\text{A.22})$$

$$\left\langle \phi'_{\frac{N}{2}}(x) \bar{\phi}'_{\frac{N}{2}}(y) \right\rangle. \quad (\text{A.23})$$

These are necessary in the evaluation of the axial anomaly in sections 4 and 5. We note that the correlation function for the Pauli-Villars field is obtained from the result for the domain-wall fermion by a simple replacement

$$\alpha_0 = \frac{1}{a_5} \begin{pmatrix} B & -C^\dagger \\ 0 & 0 \end{pmatrix} \implies \alpha_1 = \frac{1}{a_5} \begin{pmatrix} B & -C^\dagger \\ 0 & 1 \end{pmatrix}, \quad (\text{A.24})$$

$$\beta_0 = \frac{1}{a_5} \begin{pmatrix} 0 & 0 \\ C & B \end{pmatrix} \implies \beta_1 = \frac{1}{a_5} \begin{pmatrix} 1 & 0 \\ C & B \end{pmatrix}. \quad (\text{A.25})$$

For this purpose, we introduce the source terms for the $\frac{N}{2}$ -th flavor fermion in addition to those for the light N -th flavor,

$$a^4 \sum_x \left\{ \bar{J}_{\frac{N}{2}}(x) \psi_{\frac{N}{2}}(x) + \bar{\psi}_{\frac{N}{2}}(x) J_{\frac{N}{2}}(x) \right\}. \quad (\text{A.26})$$

In the upper-triangle basis of section 2, these source terms may be expressed as follows:

$$a^4 \sum_x \left\{ \bar{J}_{\frac{N}{2}}(x) \left[P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \psi''_{\frac{N}{2}}(x) + P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \psi''_{\frac{N}{2}+1}(x) \right] \right. \\ \left. + \bar{\psi}''_{\frac{N}{2}}(x) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} J_{\frac{N}{2}}(x) \right\}. \quad (\text{A.27})$$

After the integration of the first $\frac{N}{2} - 1$ flavors, the terms which contains the $\frac{N}{2}$ -th flavor are found as follows:

$$\begin{aligned}
& \left[\bar{J}(x) P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \bar{\psi}_N''(x) \beta_0 \psi_{\frac{N}{2}}''(x) \right] (-\alpha^{-1} \beta)^{\frac{N}{2}-1} \psi_{\frac{N}{2}}''(x) \\
& + \bar{\psi}_{\frac{N}{2}}''(x) \alpha \psi_{\frac{N}{2}}''(x) \\
& + \bar{\psi}_{\frac{N}{2}}''(x) \beta \psi_{\frac{N}{2}+1}''(x) \\
& + \bar{J}_{\frac{N}{2}}(x) P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \psi_{\frac{N}{2}}''(x) + \bar{\psi}_{\frac{N}{2}}''(x) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} J_{\frac{N}{2}}(x).
\end{aligned} \tag{A.28}$$

Integrating the $\frac{N}{2}$ -th flavor, we obtain

$$\begin{aligned}
& \left\{ \left[\bar{J}(x) P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \bar{\psi}_N''(x) \beta_0 \right] (-\alpha^{-1} \beta)^{\frac{N}{2}-1} \right. \\
& \quad \left. + \bar{J}_{\frac{N}{2}}(x) P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\} (-\alpha^{-1}) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} J_{\frac{N}{2}}(x) \\
& + \left\{ \left[\bar{J}(x) P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \bar{\psi}_N''(x) \beta_0 \right] (-\alpha^{-1} \beta)^{\frac{N}{2}} \right. \\
& \quad \left. + \bar{J}_{\frac{N}{2}}(x) \left[P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (-\alpha^{-1} \beta) + P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right] \right\} \psi_{\frac{N}{2}+1}''(x) \\
& \quad + \bar{\psi}_{\frac{N}{2}+1}''(x) \alpha \psi_{\frac{N}{2}+1}''(x) \\
& \quad + \bar{\psi}_{\frac{N}{2}+1}''(x) \beta \psi_{\frac{N}{2}+2}''(x).
\end{aligned} \tag{A.29}$$

Further integration up to the $N - 1$ -th flavor results in

$$\begin{aligned}
& \bar{J}(x) \left[P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (-\alpha^{-1} \beta)^{N-1} + P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right] \psi_N''(x) \\
& + \bar{\psi}_N''(x) \left[\alpha_0 + \beta_0 (-\alpha^{-1} \beta)^{N-1} \right] \psi_N''(x) + \bar{\psi}_N''(x) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} J(x) \\
& + \bar{J}_{\frac{N}{2}}(x) \left[P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (-\alpha^{-1} \beta) + P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right] (-\alpha^{-1} \beta)^{\frac{N}{2}-1} \psi_N''(x) \\
& \quad + \bar{\psi}_N''(x) \beta_0 (-\alpha^{-1} \beta)^{\frac{N}{2}-1} (-\alpha^{-1}) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} J_{\frac{N}{2}}(x) \\
& + \left\{ \bar{J}(x) P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (-\alpha^{-1} \beta)^{\frac{N}{2}-1} + \bar{J}_{\frac{N}{2}}(x) P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right\} \times \\
& \quad (-\alpha^{-1}) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} J_{\frac{N}{2}}(x).
\end{aligned} \tag{A.30}$$

Noting Eq. (2.38);

$$\begin{aligned}\alpha_0 \alpha^{-1} &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \\ \beta_0 \beta^{-1} &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},\end{aligned}\quad (\text{A.31})$$

$$(-\beta \alpha^{-1}) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \gamma_5 e^{a_5 \tilde{H}} \gamma_5 \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (\text{A.32})$$

we can evaluate the factor of the first term as follows:

$$\begin{aligned}&\left[P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (-\alpha^{-1} \beta) + P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right] (-\alpha^{-1} \beta)^{\frac{N}{2}-1} \alpha^{-1} \\ &= a_5 e^{\frac{N}{2} a_5 \tilde{H}} (-\gamma_5) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.\end{aligned}\quad (\text{A.33})$$

Combining with Eq. (2.39);

$$\begin{aligned}&\left[P_L \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (-\alpha^{-1} \beta)^{N-1} + P_R \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right] \alpha^{-1} \\ &= -a_5 \left[P_R + P_L e^{N a_5 \tilde{H}} \right] (-\gamma_5) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},\end{aligned}\quad (\text{A.34})$$

$$\begin{aligned}&[\alpha_0 + \beta_0 (-\alpha^{-1} \beta)^{N-1}] \alpha^{-1} \\ &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \left[P_L + P_R e^{N a_5 \tilde{H}} \right] (-\gamma_5) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},\end{aligned}\quad (\text{A.35})$$

we obtain

$$\begin{aligned}\left\langle \psi'_{\frac{N}{2}}(x) \bar{q}(y) \right\rangle &= \frac{a_5}{a^4} e^{\frac{N}{2} a_5 \tilde{H}} \frac{1}{P_L + P_R e^{N a_5 \tilde{H}}} \\ &= \frac{a_5}{a^4} \frac{1}{\cosh \frac{N}{2} a_5 \tilde{H}} \frac{1}{1 + \gamma_5 \tanh \frac{N}{2} a_5 \tilde{H}}.\end{aligned}\quad (\text{A.36})$$

As to the correlation function of the $\frac{N}{2}$ -th flavor, we obtain

$$\begin{aligned}\left\langle \psi'_{\frac{N}{2}}(x) \bar{\psi}'_{\frac{N}{2}}(y) \right\rangle &= \frac{a_5}{a^4} \left(-P_R + e^{\frac{N}{2} a_5 \tilde{H}} \frac{1}{P_L + P_R e^{N a_5 \tilde{H}}} P_R e^{\frac{N}{2} a_5 \tilde{H}} \gamma_5 \right) \\ &= \frac{a_5}{a^4} \left(-P_R + \frac{1}{P_L e^{-\frac{N}{2} a_5 \tilde{H}} + P_R e^{\frac{N}{2} a_5 \tilde{H}}} P_R e^{\frac{N}{2} a_5 \tilde{H}} \gamma_5 \right) \\ &= \frac{a_5}{a^4} \left(-P_L - \frac{1}{P_L e^{-\frac{N}{2} a_5 \tilde{H}} + P_R e^{\frac{N}{2} a_5 \tilde{H}}} P_L e^{-\frac{N}{2} a_5 \tilde{H}} \gamma_5 \right).\end{aligned}\quad (\text{A.37})$$

By averaging the last two expressions of Eq. (A.37), we have

$$\begin{aligned} \left\langle \psi'_{\frac{N}{2}}(x) \bar{\psi}'_{\frac{N}{2}}(y) \right\rangle &= \frac{a_5}{a^4} \left\{ -\frac{1}{2} + \frac{1}{2} \frac{1}{\cosh \frac{N}{2} a_5 \tilde{H}} \times \right. \\ &\quad \frac{1}{1 + \gamma_5 \tanh \frac{N}{2} a_5 \tilde{H}} \gamma_5 \left(1 + \gamma_5 \tanh \frac{N}{2} a_5 \tilde{H} \right) \times \\ &\quad \left. \cosh \frac{N}{2} a_5 \tilde{H} \gamma_5 \right\}. \end{aligned} \quad (\text{A.38})$$

We then remind the following identity,

$$\gamma_5 D_N = -D_N \gamma_5 (1 - a R D_N) + \Delta_N, \quad (\text{A.39})$$

where

$$D_N = \frac{1}{2a} \left(1 + \gamma_5 \tanh \frac{N}{2} a_5 \tilde{H} \right) \quad (\text{A.40})$$

$$\gamma_5 (1 - a R D_N) = -\tanh \frac{N}{2} a_5 \tilde{H}, \quad (\text{A.41})$$

and

$$\Delta_N = \gamma_5 \frac{1}{a} \frac{1}{\cosh^2 \frac{N}{2} a_5 \tilde{H}}. \quad (\text{A.42})$$

Then we finally obtain the following expression

$$\begin{aligned} \left\langle \psi'_{\frac{N}{2}}(x) \bar{\psi}'_{\frac{N}{2}}(y) \right\rangle &= \frac{a_5}{a^4} \left(-1 + \frac{1}{2} \gamma_5 a R D_N \gamma_5 \right. \\ &\quad \left. + \frac{1}{2a} \frac{1}{\cosh \frac{N}{2} a_5 \tilde{H}} \frac{1}{D_N} \gamma_5 \frac{1}{\cosh \frac{N}{2} a_5 \tilde{H}} \gamma_5 \right). \end{aligned} \quad (\text{A.43})$$

In the case of the Pauli-Villars field, through the replacement of Eq. (A.24), we obtain

$$\begin{aligned} \alpha_1 \alpha^{-1} &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (-\gamma_5) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \\ \beta_1 \beta^{-1} &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} (+\gamma_5) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \end{aligned} \quad (\text{A.44})$$

and

$$\begin{aligned} &\left[\alpha_1 + \beta_1 (-\alpha^{-1} \beta)^{N-1} \right] \alpha^{-1} \\ &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \left[1 + e^{N a_5 \tilde{H}} \right] (-\gamma_5) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \end{aligned} \quad (\text{A.45})$$

instead of Eqs. (A.31) and (A.35). Then we can evaluate the correlation function as

$$\begin{aligned}
\left\langle \phi'_{\frac{N}{2}}(x) \bar{\phi}'_{\frac{N}{2}}(y) \right\rangle &= \frac{a_5}{a^4} \left(-P_R + e^{\frac{N}{2}a_5\tilde{H}} \frac{1}{1 + e^{Na_5\tilde{H}}} e^{\frac{N}{2}a_5\tilde{H}} \gamma_5 \right) \\
&= \frac{a_5}{a^4} \left(-\frac{1}{2} + \frac{1}{2} \tanh \frac{N}{2} a_5 \tilde{H} \gamma_5 \right) \\
&= -\gamma_5 \frac{a_5}{a^4} (1 - aD_N) \gamma_5.
\end{aligned} \tag{A.46}$$

References

- [1] H. Neuberger, Phys. Lett. **B417** (1998) 141; Phys. Lett. **B427** (1998) 353.
- [2] P. Hasenfratz, Nucl. Phys. **B**(Proc. Suppl.) **63** (98) 53.
- [3] P. Hernández, K. Jansen and M. Lüscher, [hep-lat/9808010](#).
- [4] P. H. Ginsparg and K. G. Wilson, Phys. Rev. **D25** (1982) 2649.
- [5] M. Lüscher, Phys. Lett. **B428** (1998) 342.
- [6] R. Narayanan and H. Neuberger, Nucl. Phys. **B412** (1994) 574; Phys. Rev. Lett. **71** (1993) 3251; Nucl. Phys. **B443** (1995) 305.
- [7] H. Neuberger, Phys. Rev. Lett. **81** (1998) 4060;
“Minimizing storage in implementations of the overlap lattice-Dirac operator”, [hep-lat/9811019](#); “The overlap lattice Dirac operator and dynamical fermions”, [hep-lat/9901003](#).
- [8] R.G. Edwards, Urs M. Heller and R. Narayanan, “A Study of Practical Implementations of the Overlap-Dirac Operator in Four Dimensions”, [hep-lat/9807017](#); “A study of chiral symmetry in quenched QCD using the Overlap-Dirac operator”, [hep-lat/9811030](#).
- [9] T.-W. Chiu, Phys. Rev. **D58** (1998) 074511; “GW fermion propagators and chiral condensate”, [hep-lat/9810052](#).
- [10] A. Borici, On the Neuberger overlap operator, [hep-lat/9810064](#).
- [11] D. B. Kaplan, Phys. Lett. **B288** (1992) 342.
- [12] Y. Shamir, Nucl. Phys. **B406** (1993) 90.

- [13] V. Furman and Y. Shamir, Nucl. Phys. **B439** (1995) 54.
- [14] H. Neuberger, Phys. Rev. **D57** (1998) 5417.
- [15] P. Vranas, Phys. Rev. **D57** (1998) 1415.
- [16] T. Blum and A. Soni, Phys. Rev. **D56** (1997) 174; Phys. Rev. Lett. **79** (1997) 3595; Update on lattice QCD with domain wall quarks, [hep-lat/9712004](#).
- [17] S. Chandrasekharan, D. Chen, N. Christ, W. Lee, R. Mawhinney, P. Vranas, Anomalous Chiral Symmetry Breaking above the QCD Phase Transition, [hep-lat/9807018](#). P. Chen, N. Christ, G. Fleming, A. Kaehler, C. Malureanu, R. Mawhinney, C. Sui, P. Vranas, Y. Zhestkov Domain wall fermion zero modes on classical topological backgrounds, [hep-lat/9807029](#); P. Chen, N. Christ, G. Fleming, A. Kaehler, C. Malureanu, R. Mawhinney, G. Siebert, C. Sui, P. Vranas, Y. Zhestkov, Dynamical QCD thermodynamics with domain wall fermions, [hep-lat/9809159](#); The domain wall fermion chiral condensate in quenched QCD, [hep-lat/9811013](#); Quenched QCD with domain wall fermions, [hep-lat/9811026](#);
- [18] T. Blum, A. Soni, M. Wingate, Light quark masses using domain wall fermions, [hep-lat/9809065](#).
- [19] J.-F. Lagae, D. K. Sinclair, Domain wall fermions at finite temperature, [hep-lat/9809134](#).
- [20] S. Aoki, Y. Taniguchi, One loop calculation in lattice QCD with domain-wall quarks, [hep-lat/9711004](#); One loop renormalization for the axial Ward-Takahashi identity in Domain-wall QCD, [hep-lat/9811007](#).
- [21] Y. Kikukawa, H. Neuberger and A. Yamada, Nucl. Phys. **B526** (1998) 572.
- [22] T. Blum, Domain wall fermions in vector gauge theories, Plenary review talk at LATTICE98, [hep-lat/9810017](#).
- [23] P. Vranas, Phys. Rev. **D57** (1998) 1415.
- [24] T.-W. Chiu and S.V. Zenkin, Solutions of the Ginsparg-Wilson Relation, [hep-lat/9806019](#); T.-W. Chiu, C.-W. Wang and S.V. Zenkin, Chiral structure of the solutions of the Ginsparg-Wilson relation

- Phys.Lett. B438 (1998) 321; T.-W. Chiu, The axial anomaly of Ginsparg-Wilson fermion, [hep-lat/9809013](#).
- [25] F. Niedermayer, plenary talk given by F. Niedermayer at the International Symposium on Lattice Field Theory, Boulder, July 13-18, 1998.
- [26] Y. Shamir, Nucl. Phys. B417 (1994) 167.
- [27] P. Hasenfratz, V. Laliena and F. Niedermayer, Phys.Lett. B427 (1998) 125.
- [28] Y. Kikukawa and A. Yamada, “Weak coupling expansion of massless QCD with a Ginsparg-Wilson fermion and axial U(1) anomaly”, [hep-lat/9806013](#), to appear in Phys. Lett. B.
 K. Fujikawa, A continuum limit of the chiral Jacobian in lattice gauge theory, [hep-th/9811235](#), To be published in Nuclear Physics B.
 D.H. Adams, Axial anomaly and topological charge in lattice gauge theory with overlap-Dirac operator, [hep-lat/9812003](#).
 H. Suzuki, Simple Evaluation of Chiral Jacobian with Overlap Dirac Operator, [hep-th/9812019](#).
- [29] Y. Kikukawa and A. Yamada, Axial vector current of exact chiral symmetry on the lattice, [hep-lat/9808026](#), to appear in Nucl. Phys. B.